


Procedure date: _____ Procedure Time: _____ If you need to reschedule, call: <input type="checkbox"/> Charlynn - (912)777-9032 <input type="checkbox"/> Cynthia - (912)790-2649 <input type="checkbox"/> Briana - (912)790-2645 <input type="checkbox"/> Richard - (912)790-2685 <input type="checkbox"/> Heather - (912)790-2827 Location to report to: o The Endoscopy Center – (912)303-4200 / (1139 Lexington Avenue) o Bluffton Okatie Surgery Center–(912)303-4200 / (40 Okatie Ctr Blvd S Ste 125, Okatie, SC) o Memorial – (912)350-7832 / (Memorial Outpatient & Wellness Center) o St. Joseph’s Hospital – (912)819-2171 / (Day Surgery/Imaging Emergency Entrance, 2 nd Floor) o Candler Hospital – (912)819-6800 / (Professional Building)	Check in Time: _____ Physician: _____ FOLLOW UP APPOINTMENT is scheduled for: Date: _____ Check in Time: _____ Appointment Time: _____ Physician: _____ Mid-level Provider: _____ Location to report to: o 1139 Lexington Avenue, Savannah—(912) 303-4200 o 140 Traders Way, Pooler o 40 Okatie Ctr Blvd S Ste 210, Okatie, SC
---	---

COLONOSCOPY – procedure performed to help identify and/or correct a problem in the colon. You will be sedated before an endoscope is inserted through the anus and gently moved around the bends of the colon. The procedure usually takes 30-60 minutes.

- Do NOT Stop taking:** Aggrenox Brilinta Coumadin Effient Eliquis Plavix Pradaxa _____
 STOP taking: Aggrenox Brilinta Coumadin Effient Eliquis Plavix Pradaxa _____ for _____ days before procedure

Have you signed up for your patient portal yet? If not, please visit our website www.savannahgi.com to sign up and you can also find additional information such as: Colonoscopy prep instructions, FAQ’s and additional services provided.

****To ensure that you are prepped correctly, please ONLY follow these patient instructions - DO NOT follow instructions from the pharmacist or the prep ox****

<p><u>7 DAYS BEFORE PROCEDURE</u></p> <p> STOP ALL iron tablets and fiber supplements: (Metamucil, Citrucel, Fiber-con, Benefiber, Konsyl, etc) along with corn, beans, nuts and seeds.</p> <p>Make sure you have your bowel preparation.</p> <p>Make arrangements for a responsible adult to come with you and REMAIN in The Endoscopy Center until you are discharged. Your responsible adult must drive you home and be present with you the remainder of day.</p> <p>There are no exceptions. You are not allowed to take a bus, taxi or leave The Endoscopy Center ALONE. If you do not have a responsible adult with you to take you home, your exam cannot be done and will be cancelled.</p>	<p><u>2 DAYS BEFORE PROCEDURE</u> If unable to keep your procedure appointment, please call 912-303-4200 to cancel. For patients who do not show for a procedure or do not cancel 2 days prior to their scheduled procedure time, a \$150 fee will be charged.</p> <p><u>1 DAY BEFORE THE PROCEDURE – Date:</u> _____ Begin a clear liquid diet the morning prior to your procedure and continue throughout the day. Adequate hydration is very important.</p> <p>This consists of coffee and/or tea with artificial sweeteners ONLY – (NO milk, cream, or sugar). Other choices are: water, diet sodas, beef and/or chicken broth, sugar free Jell-O, and popsicles.</p> <p>AVOID red or purple food coloring.</p> <p>NO SOLID FOODS</p> <p>If you are diabetic and experiencing symptoms of low blood sugar, you may take small sips of white grape juice.</p> <p>Empty 1 Pouch A and 1 Pouch B into the disposable container. Add lukewarm drinking water to the top line of the container. Mix to dissolve. You may refrigerate prior to drinking if preferred. The solution must be consumed within 24 hours of mixing.</p>	<p><u>5:00pm THE DAY BEFORE PROCEDURE -</u> The Moviprep container is divided by 4 marks.</p> <p>Every 20 minutes, DRINK the solution down to the next mark (approximately 8 oz.) until 1 liter is completed.</p> <p>Drink 16 oz. of clear liquids of your choice after drinking the prep solution.</p> <p>You may continue to drink clear liquids to maintain hydration.</p> <p>Alcohol-free baby wipes or Vaseline may help ease skin irritation. Over the counter hydrocortisone creams or Tucks may be used if desired.</p> <p>You may pre-mix your 2nd dose and refrigerate if preferred. You will take the 2nd dose 6 hours prior to your procedure.</p>	<p><u>6 HOURS BEFORE PROCEDURE – Time:</u> The Moviprep container is divided by 4 marks. Every 20 minutes, DRINK the solution down to the next mark (approximately 8 oz) until 1 liter is completed. Drink 16 oz. of clear liquids of your choice after drinking the prep solution.</p> <p><u>THE MORNING OF YOUR PROCEDURE:</u> You should only take blood pressure, heart, thyroid and/or seizure medications with a SMALL SIP of water.</p> <p><u>4 HOURS BEFORE PROCEDURE – Time:</u> Your prep must be completed 4 hours prior to your scheduled procedure. STOP drinking ALL liquids including water. Anesthesia may cancel your procedure, if you have had ANY liquids within 4 hours before your procedure.</p> <p>If you are diabetic: Do NOT take oral medication or insulin the morning of the procedure. However, if you are on an insulin pump, continue the basal rate only.</p> <p><u>Reminders:</u> NO smoking prior to anesthesia on the day of your procedure.</p> <p><i>Please bring a list of all your current medications, including any allergy or over-the-counter medications.</i> <i>Bring your current insurance card.</i> <i>Please wear comfortable, loose fitting clothing that is easy to step into.</i> <i>Please leave all jewelry and valuables at home.</i> After your procedure is completed, you will NOT be able to drive, make any business decisions or perform any potentially risky work duties, so please plan accordingly. If you have any questions or problems, please call us at 912-303-4200.</p>
---	---	---	---